

Axiomatizing Rational Power Series

S. L. Bloom

Dept. of Computer Science
Stevens Institute of Technology
Hoboken, NJ, USA

Z. Ésik*

Dept. of Computer Science
University of Szeged
Szeged, Hungary
GRLMC
Rovira i Virgili University
Tarragona, Spain

Abstract

Iteration semirings are Conway semirings satisfying Conway's group identities. We show that the semirings $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ of rational power series with coefficients in the semiring $\mathbb{N}$ of natural numbers are the free partial iteration semirings. Moreover, we characterize the semirings $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ as the free semirings in the variety of iteration semirings defined by three additional simple identities, where $\mathbb{N}_\infty$ is the completion of $\mathbb{N}$ obtained by adding a point of infinity. We also show that this latter variety coincides with the variety generated by the complete, or continuous semirings. As a consequence of these results, we obtain that the semirings $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, equipped with the sum order, are free in the class of symmetric inductive $*$ -semirings. This characterization corresponds to Kozen's axiomatization of regular languages.

1 Introduction

One of the most basic algebraic structures studied in Computer Science are the semirings $\mathbf{Reg}(\Sigma^*)$ of regular (or rational) languages over an alphabet Σ equipped with the star operation. Salomaa [32] has axiomatized these semirings of regular languages using a few simple identities and the *unique fixed point rule* asserting that if the regular language a does not contain the empty word then a^*b is the unique solution of the fixed point equation $x = ax + b$. There are several ways of expressing the empty word property using a first-order language. Probably, the simplest way is by the inequality $1 + a \neq a$. Using this, the unique fixed point rule can be formulated as the first-order axiom

$$\forall a \forall b \forall x ((1 + a \neq a \wedge ax + b = x) \Rightarrow x = a^*b).$$

Salomaa's result then amounts to the assertion that for any Σ , $\mathbf{Reg}(\Sigma^*)$ is freely generated in the class of $*$ -semirings satisfying a finite number of (simple) identities and the above axiom. We have thus a *finite* first-order axiomatization of regular languages.

Because of the extra condition on a , the unique fixed point rule is not a quasi-identity. A finite axiomatization using only quasi-identities has been first obtained by Archangelsky and Gorshkov, cf. [2]. A second, and perhaps more serious concern is that several natural $*$ -semirings which

*Partially supported by grant no. MTM2007-63422 from the Ministry of Education and Science of Spain.

satisfy all identities of regular languages are not models of the unique fixed point rule. Examples of such semirings are semirings of binary relations with the reflexive-transitive closure operation as star, since for binary relations, the equation $x = ax + b$ usually has several solutions, even if $1 + a \neq a$ (i.e., when a not reflexive). On the other hand, a^*b is least among all solutions, so that

$$\forall a \forall b \forall x (x = ax + b \Rightarrow a^*b \leq x)$$

where $a^*b \leq x$ may be viewed as abbreviation for $a^*b + x = x$. And indeed, the semirings of regular languages can be characterized as the free algebras in a quasi-variety of semirings with a star operation axiomatized by a finite set of simple identities and the above *least fixed point rule*, or the *least pre-fixed point rule*

$$\forall a \forall b \forall x (ax + b \leq x \Rightarrow a^*b \leq x).$$

This result is due to Krob [25]. In [22, 23], Kozen also required the dual of the least (pre-)fixed point rule

$$\forall a \forall b \forall x (xa + b \leq x \Rightarrow ba^* \leq x),$$

and gave a simpler proof of completeness of this system. Several other finite axiomatizations are derivable from Krob's and Kozen's systems, see [12, 13, 9].

But the largest class of algebras in which the semirings of regular languages are free is of course a variety. This variety, the class of all semirings with a star operation satisfying all identities true of regular languages, is the same as the variety generated by all $*$ -semirings of binary relations. The question whether this variety is finitely based was answered by Redko [30, 31] and Conway [15], who showed that there is no finite (first-order or equational) axiomatization. The question of finding infinite equational bases was considered in [7, 25]. The system given in Krob [25] consists of the Conway semiring identities, the identity $1^* = 1$, and Conway's group identities [15] associated with the finite (simple) groups. Conway semirings were first defined formally in [6, 8]. Conway semirings are semirings equipped with a star operation satisfying $(a+b)^* = a^*(ba^*)^*$ and $(ab)^* = a(ba^*)^*b + 1$. Conway semirings satisfying the infinite collection of group identities are called iteration semirings, cf. [18]. The terminology is due to the fact that iteration semirings are exactly the semirings which are iteration algebras, i.e., satisfy the axioms of iteration theories [8] which capture the equational properties of the fixed point operation. Thus, Krob's result characterizes the semirings of regular languages as the free iteration semirings satisfying $1^* = 1$ (which implies that sum is idempotent). Another proof of this result using iteration theories can be obtained by combining the axiomatization of regular languages from [7] and the completeness (of certain generalizations of) the group identities for iteration theories, established in [18].

In this paper, we drop the idempotence of the sum operation and consider the semirings of rational power series $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ and $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ over the semiring $\mathbb{N}$ of natural numbers and its completion $\mathbb{N}_\infty$ with a point of infinity. The star operation in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is defined only on those proper power series having 0 as the coefficient of the empty word (the empty word property), whereas the star operation in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is totally defined. We prove that $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the variety $\mathcal{V}$ of all iteration semirings satisfying the identities $1^*1^* = 1^*$, $1^*a = a1^*$ and $1^*(1^*a)^* = 1^*a^*$. This result is also of interest because $\mathcal{V}$ coincides with the variety generated by those $*$ -semirings that arise from (countably) complete or continuous semirings by defining a^* as the sum $\sum_{n \geq 0} a^n$. Moreover, we prove that $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the class of all *partial* iteration semirings. As a consequence of the equational axiomatizations, we show that $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, equipped with the the sum order, is freely generated by Σ in the class of ordered $*$ -semirings satisfying the fixed point identity $aa^* + 1 = a^*$ and the least pre-fixed point rule.

The paper is organized as follows. In Section 2 we review the notion of semirings and power series. Section 3 is devoted to (partial) Conway and iteration semirings. In Section 4 we provide a formulation of the Kleene-Schützenberger theorem for (partial) Conway semirings from [8, 11]. In the characterization of the semirings $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ as the free partial iteration semirings, in addition

to the Kleene-Schützenberger theorem, our main tool will be the commutative identities. We establish several technical results for the commutative identity in Section 5. Section 6 is devoted to proving the freeness result for the semirings $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ mentioned above. Then, in Section 7 we prove that the semirings $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ are free in the variety if iteration semirings satisfying three additional simple identities. Last, in Section 8 we characterize the semirings $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ as the free symmetric inductive $*$ -semirings, and as the free inductive $*$ -semirings satisfying an additional inequation.

2 Semirings

A *semiring* [20] is an algebra $S = (S, +, \cdot, 0, 1)$ such that $(S, +, 0)$ is a commutative monoid, where $+$ is called sum or addition, $(S, \cdot, 1)$ is a monoid, where $\cdot$ is called product or multiplication. Moreover, 0 is an absorbing element with respect to multiplication and product distributes over sum:

$$\begin{aligned} 0 \cdot a &= 0 \\ a \cdot 0 &= 0 \\ a(b + c) &= ab + ac \\ (b + c)a &= ba + ca \end{aligned}$$

for all $a, b, c \in S$. A semiring S is called *idempotent* if

$$a + a = a$$

for all $a \in S$. A morphism of semirings preserves the sum and product operations and the constants 0 and 1 . Since semirings are defined by identities, the class of all semirings is a variety (see e.g., [21]) as is the class of all idempotent semirings.

An important example of a semiring is the semiring $\mathbb{N} = (\mathbb{N}, +, \cdot, 0, 1)$ of natural numbers equipped with the usual sum and product operations. An important example of an idempotent semiring is the boolean semiring $\mathbb{B}$ whose underlying set is $\{0, 1\}$ and whose sum and product operations are the operations $\vee$ and $\wedge$, i.e., disjunction and conjunction. Actually $\mathbb{N}$ and $\mathbb{B}$ are respectively the initial semiring and the initial idempotent semiring. The semiring $\mathbb{N}_\infty$ is defined on the set $\mathbb{N} \cup \{\infty\}$ so that it contains $\mathbb{N}$ as a subsemiring and $n + \infty = \infty + n = \infty$ and $m\infty = \infty m = \infty$ for all $n, m \in \mathbb{N} \cup \{\infty\}$, $m \neq 0$.

We describe two constructions on semirings. For more information on semirings, the reader is referred to Golan's book [20].

The first construction is that of matrix semirings. When S is a semiring, then for each $n \geq 0$ the set $S^{n \times n}$ of all $n \times n$ matrices over S is also a semiring. The sum operation is defined pointwise and product is the usual matrix product. The constants are the matrix 0_n all of whose entries are 0 (often denoted just 0), and the diagonal matrix E_n whose diagonal entries are all 1 . In addition to square matrices, we will also consider more general rectangular matrices with the usual definition of sum and product. (Rectangular matrices over S form a semiadditive category that can canonically be assigned to S but we will avoid using categorical notions.) When ρ is a function $\{1, \dots, n\} \rightarrow \{1, \dots, p\}$, for some $n, p \geq 0$, there is a corresponding $n \times p$ matrix over each semiring: it is a 0-1 matrix with a 1 on the (i, j) th position exactly when $i\rho = j$. Such matrices will be called *functional*. A *permutation matrix* is a functional matrix that corresponds to a permutation.

The second construction is that of power series and polynomial semirings, cf. [5]. Suppose that S is a semiring and Σ is a set. Let Σ^* denote the free monoid of all words over Σ including the empty

word ϵ . A *power series* over S (in the noncommuting letters in Σ) is a function $s : \Sigma^* \rightarrow S$. It is a common practice to represent a power series s as a formal sum $\sum_{w \in \Sigma^*} (s, w)w$, where the *coefficient* (s, w) is ws , the value of s on the word w . The *support* of a series s is the set $\text{supp}(s) = \{w : (s, w) \neq 0\}$. When $\text{supp}(s)$ is finite, s is called a *polynomial*. We let $S\langle\langle\Sigma^*\rangle\rangle$ and $S\langle\Sigma^*\rangle$ respectively denote the collection of all power series and polynomials over S in the letters Σ . More generally, when $S' \subseteq S$, we let $S'\langle\langle\Sigma^*\rangle\rangle$ denote the set of all power series in $S\langle\langle\Sigma^*\rangle\rangle$ all of whose coefficients are in S' . The set of polynomials $S'\langle\Sigma^*\rangle$ is defined in the same way. We denote by $S'\langle\Sigma\rangle$ (no star) the collection of those polynomials in $S'\langle\Sigma^*\rangle$ which are linear combinations over Σ .

We define the sum $s + s'$ and product $s \cdot s'$ of two series $s, s' \in S\langle\langle\Sigma^*\rangle\rangle$ as follows. For all $w \in \Sigma^*$,

$$\begin{aligned} (s + s', w) &= (s, w) + (s', w) \\ (s \cdot s', w) &= \sum_{uu' = w} (s, u)(s', u'). \end{aligned}$$

We may identify any element $s \in S$ with the series, in fact polynomial, which maps ϵ to s and all other elements of Σ^* to 0. In particular, 0 and 1 may be viewed as polynomials. It is well-known that equipped with the above operations and constants, $S\langle\langle\Sigma^*\rangle\rangle$ is a semiring which contains $S\langle\Sigma^*\rangle$ as a subsemiring.

Note that $\mathbb{B}\langle\langle\Sigma^*\rangle\rangle$ is isomorphic to the semiring of languages in Σ^* , where addition corresponds to set union and multiplication to concatenation. An isomorphism maps each series in $\mathbb{B}\langle\langle\Sigma^*\rangle\rangle$ to its support, and the inverse of this isomorphism maps each language $L \subseteq \Sigma^*$ to its *characteristic series* s defined by $(s, w) = 1$ if $w \in L$ and $(s, w) = 0$, otherwise.

The following fact is well-known.

THEOREM 2.1 *Given any semirings S, S' , any semiring morphism $h_S : S \rightarrow S'$ and any function $h : \Sigma \rightarrow S'$ such that*

$$(sh_S)(ah) = (ah)(sh_S) \tag{1}$$

for all $a \in \Sigma$ and $s \in S$, there is a unique semiring morphism $h^\sharp : S\langle\Sigma^*\rangle \rightarrow S'$ which extends both h_S and h .

The condition (1) means that for any $s \in S$ and letter $a \in \Sigma$, sh_S commutes with ah . In particular, since $\mathbb{N}$ is initial, and since when $S = \mathbb{N}$ the condition (1) holds automatically, we obtain that any map $\Sigma \rightarrow S'$ into a semiring S' extends to a unique semiring morphism $\mathbb{N}\langle\Sigma^*\rangle \rightarrow S'$, i.e., the polynomial semiring $\mathbb{N}\langle\Sigma^*\rangle$ is freely generated by Σ in the variety of semirings. In the same way, $\mathbb{B}\langle\Sigma^*\rangle$ is freely generated by Σ in the variety of idempotent semirings.

3 Conway and iteration semirings

In this section, we review the notions of (partial) Conway semiring and iteration semiring. The notions and facts presented here will be used in the freeness results.

The definition of Conway semirings involves two important identities of regular languages. They appear implicitly in Conway [15] and were first defined explicitly in [6, 8]. Partial Conway semirings appear in [11]. Recall that an *ideal* of a semiring S is a set $I \subseteq S$ which contains 0 and satisfies $I + I \subseteq I$ and $SI \cup IS \subseteq I$.

This section is based on [8] and [11].

DEFINITION 3.1 A partial $*$ -semiring is a semiring equipped with a partially defined star operation $a \mapsto a^*$ whose domain $D(S)$ is an ideal. A partial Conway semiring is a $*$ -semiring S satisfying the sum star and product star identities:

1. Sum star identity:

$$(a + b)^* = a^*(ba^*)^* \quad (2)$$

for all $a, b \in D(S)$.

2. Product star identity:

$$(ab)^* = 1 + a(ba)^*b, \quad (3)$$

for all $a, b \in S$ such that $a \in D(S)$ or $b \in D(S)$.

A $*$ -semiring is a partial $*$ -semiring S with $D(S) = S$, i.e., the star operation is totally defined. A Conway semiring is a partial Conway semiring which is a $*$ -semiring. Morphisms $h : S \rightarrow S'$ of (partial) $*$ -semirings and (partial) Conway semirings preserve the ideal and star operation: if $a \in D(S)$ then $ah \in D(S')$ and $a^*h = (ah)^*$.

Note that in any partial Conway semiring S ,

$$aa^* + 1 = a^* \quad (4)$$

$$a^*a + 1 = a^* \quad (5)$$

$$0^* = 1 \quad (6)$$

for all $a \in D(S)$. Moreover, for all $a, b \in S$ with $a \in D(S)$ or $b \in D(S)$,

$$(ab)^*a = a(ba)^*. \quad (7)$$

It then follows that also

$$(a + b)^* = (a^*b)^*a^* \quad (8)$$

for all $a, b \in D(S)$, which can be used instead of (2) in the definition of partial Conway semirings. By (4) and (5), for any a, b in a partial Conway semiring S , if $a \in D(S)$ then a^*b is a solution of the equation $x = ax + b$ and ba^* is a solution of $x = xa + b$. In particular, $a^+ = aa^* = a^*a$ is a solution of both $x = ax + a$ and $x = xa + a$.

An important feature of (partial) Conway semirings is that square matrices over Conway semirings also form (partial) Conway semirings.

DEFINITION 3.2 Suppose that S is a partial Conway semiring. We turn the semirings $S^{k \times k}$, $k \geq 0$ into partial $*$ -semirings. Note that $D(S)^{k \times k}$, the collection of all $k \times k$ matrices all of whose entries are in $D(S)$ is an ideal of $S^{k \times k}$. The star operation will be defined on this ideal. When $k = 0$, $S^{k \times k}$ is trivial as is the definition of star. When $k = 1$, we use the star operation on S . Assuming that $k > 1$ we write $k = n + 1$. For a matrix $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ define

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^* = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \quad (9)$$

where $A \in D(S)^{n \times n}$, $B \in D(S)^{n \times 1}$, $C \in D(S)^{1 \times n}$, and $D \in D(S)^{1 \times 1}$, and where

$$\begin{aligned} \alpha &= (A + BD^*C)^* & \beta &= \alpha BD^* \\ \gamma &= \delta CA^* & \delta &= (D + CA^*B)^*. \end{aligned}$$

PROPOSITION 3.3 *When S is a (partial) Conway semiring, so is each $S^{n \times n}$. Moreover, the matrix star identity (9) holds for all matrices $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ with $A \in D(S)^{n \times n}$, $B \in D(S)^{n \times m}$, $C \in D(S)^{m \times n}$, and $D \in D(S)^{m \times m}$, all $n, m \geq 0$.*

In fact, $(AB)^* = A(BA)^*B + E_n$ holds for all rectangular matrices $A \in D(S)^{n \times m}$ and $B \in D(S)^{m \times n}$.

For later use we note that the following *permutation identity* holds in all (partial) Conway semirings.

PROPOSITION 3.4 *When S is a partial Conway semiring, $A \in D(S)^{n \times n}$ and π is an $n \times n$ permutation matrix with transpose π^T , then $(\pi A \pi^T)^* = \pi A^* \pi^T$.*

Following Conway [15], we associate an identity in (partial) Conway semirings with each finite group. Let G be a finite group of order n . Without loss of generality we may assume that the elements of G are the integers $1, \dots, n$. Moreover, because the permutation identity holds in all (partial) Conway semirings, without loss of generality we may fix a sequencing of the elements and assume that 1 is the unit element of G .

DEFINITION 3.5 *We say that the group identity associated with a finite group G of order n holds in a partial Conway semiring S if*

$$e_1 M_G^* u_n = (a_1 + \dots + a_n)^* \quad (10)$$

holds, where $a_1, \dots, a_n$ are arbitrary elements in $D(S)$, and where M_G is the $n \times n$ matrix whose (i, j) th entry is a_{i-1j} , for all $1 \leq i, j \leq n$, and e_1 is the $1 \times n$ 0-1 matrix whose first entry is 1 and whose other entries are 0, finally u_n is the $n \times 1$ matrix all of whose entries are 1.

Equation (10) asserts that the sum of the entries of the first row of M_G^* is $(a_1 + \dots + a_n)^*$. For example, the group identity associated with the group of order 2 is

$$\begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} a_1 & a_2 \\ a_2 & a_1 \end{pmatrix}^* \begin{pmatrix} 1 \\ 1 \end{pmatrix} = (a_1 + a_2)^*,$$

which by the matrix star identity can be written as

$$(a_1 + a_2 a_1^* a_2)^* (1 + a_2 a_1^*) = (a_1 + a_2)^*.$$

(It is known that in Conway semirings, this identity is further equivalent to $(a^2)^* (1 + a) = a^*$.)

DEFINITION 3.6 *We say that a Conway semiring S is an iteration semiring if it satisfies all group identities. We say that a partial Conway semiring S is a partial iteration semiring if it satisfies all group identities (10) where $a_1, \dots, a_n$ range over $D(S)$. A morphism of (partial) iteration semirings is a (partial) Conway semiring morphism.*

We end this section by recalling from [8, 11] that power series semirings are (partial) iteration semirings. Suppose that S is a semiring and Σ is a set, and consider the semiring $S\langle\langle \Sigma^* \rangle\rangle$. A series $s \in S\langle\langle \Sigma^* \rangle\rangle$ is called *proper* [5] if $(s, \epsilon) = 0$. It is clear that the proper series form an ideal of $S\langle\langle \Sigma^* \rangle\rangle$. It is well-known (see e.g. [5]) that when s is proper and r is any series, there is a unique series that solves the fixed point equation $x = sx + r$, and that this solution is $s^* r$, where s^* is the unique solution of $y = sy + 1$.

PROPOSITION 3.7 *For any semiring S , the power series semiring $S\langle\langle\Sigma^*\rangle\rangle$, equipped with the star operation defined on proper series, is a partial iteration semiring.*

When S is a $*$ -semiring, it is possible to turn star into a total operation. Given a series $s \in S\langle\langle\Sigma^*\rangle\rangle$, it can be written in a unique way as $s = s_0 + r$, where $s_0 \in S$ and r is proper. Since s_0 is in S and S has a star operation, s_0^* is defined. We define $s^* = (s_0^*r)^*s_0^*$, where $(s_0^*r)^*$ is the unique solution of the equation $x = (s_0^*r)x + 1$ as before. The following fact is a special case of a more general result proved in [6, 8].

PROPOSITION 3.8 *When S is an iteration semiring, so is $S\langle\langle\Sigma^*\rangle\rangle$.*

4 The Kleene-Schützenberger theorem

Let S denote a semiring, let Σ denote a set, and consider the power series semiring $S\langle\langle\Sigma^*\rangle\rangle$ which is a partial iteration semiring (or an iteration semiring, if S is). As usual, we identify each letter in Σ and each element of S with a series. We call a series s in $S\langle\langle\Sigma^*\rangle\rangle$ *rational* if s belongs to the least partial iteration subsemiring of $S\langle\langle\Sigma^*\rangle\rangle$ containing $S \cup \Sigma$, i.e. when s is contained in the least subsemiring of $S\langle\langle\Sigma^*\rangle\rangle$ containing $S \cup \Sigma$ closed under the star operation. We let $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ denote the partial iteration semiring of all rational power series in $S\langle\langle\Sigma^*\rangle\rangle$. The Kleene-Schützenberger theorem [5] equates rational power series with the power series recognizable by (weighted) automata. For later use, below we give a general definition of automata applicable to all partial Conway semirings, see [11] and [8].

DEFINITION 4.1 *Let S be a partial Conway semiring and suppose that S_0 is a subsemiring of S and Σ is a subset of $D(S)$. An automaton in S over (S_0, Σ) is a triplet $\mathbf{A} = (\alpha, A, \beta)$, where for some integer n , $\alpha \in S_0^{1 \times n}$, $\beta \in S_0^{n \times 1}$, and $A \in (S_0\Sigma)^{n \times n}$, where $S_0\Sigma$ is the set of all linear combinations of the elements of Σ with coefficients in S_0 . The integer n is called the dimension of $\mathbf{A}$. The behavior of $\mathbf{A}$ is $|\mathbf{A}| = \alpha A^* \beta$.*

Thus, when the partial Conway semiring is $S\langle\langle\Sigma^*\rangle\rangle$, where S is a semiring, S_0 is S and Σ is the collection of power series corresponding to the letters in Σ , we obtain the usual notion of a (weighted) automaton. We let $S^{\text{rec}}\langle\langle\Sigma^*\rangle\rangle$ denote the collection of all power series which are behaviors of such automata. The Kleene-Schützenberger theorem is:

THEOREM 4.2 $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle = S^{\text{rec}}\langle\langle\Sigma^*\rangle\rangle$.

For a proof, see [11]. Below we will call an automaton (over (S, Σ)) in $S\langle\langle\Sigma^*\rangle\rangle$ also an automaton in $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$. When $\mathbf{A} = (\alpha, A, \beta)$ is an automaton in $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ and h is a function $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle \rightarrow S'$ into a partial iteration semiring which is a semiring morphism on S , maps Σ into $D(S')$ and preserves linear combinations in $S\langle\Sigma\rangle$, then $\mathbf{A}h = ((\alpha h), (Ah), (\beta h))$ is an automaton in S' (over $(Sh, \Sigma h)$).

For later use we also give the following result from [11].

THEOREM 4.3 *Suppose that S is a semiring and Σ is a set, so that $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is a partial iteration semiring. Suppose that S' is a partial iteration semiring and h is a function $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle \rightarrow S'$. Then h is a morphism of partial iteration semirings iff the restriction of h onto S is a semiring morphism, h maps Σ to $D(S')$ and preserves linear combinations in $S\langle\Sigma\rangle$; moreover, h preserves the behavior of automata, so that $|\mathbf{A}|h = |\mathbf{A}h|$ for all automata $\mathbf{A}$ in $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$.*

5 The commutative identity

In the proof of our results, we will deduce the equality $A^*\rho = \rho B^*$ from the equality $A\rho = \rho B$, where A is an $m \times m$, B is an $n \times n$ matrix over a partial iteration semiring, and ρ is an $m \times n$ functional matrix. The commutative identity, defined below, is a generalization of the group identities which holds in all (partial) iteration semirings. The commutative identity allows us to infer the implication above, under certain conditions. The commutative identity was introduced for $*$ -semirings in [8] but its origins in iteration theories go back to [17]. See also [18]. This section is rather technical and all proofs may be skipped at first reading.

In order to illustrate the commutative identity and its use, consider the following situation. Assume that A , B and ρ are as above, but for simplicity assume that ρ as a function is surjective and monotone, collapsing the first m_1 integers to 1, the next m_2 integers to 2 etc. Then write A as a block matrix $(A_{ij})_{ij}$, where each A_{ij} is a $m_i \times m_j$ matrix for all $i, j = 1, \dots, n$. The condition that $A\rho = \rho B$ means that each row sum of any A_{ij} is b_{ij} , the (i, j) th entry of matrix B . Similarly, $A^*\rho = \rho B^*$ means that A^* can be written as a matrix of blocks of size $m_i \times m_j$, $i, j = 1, \dots, n$, and for each i and j , each row sum of the (i, j) th block is equal to the (i, j) th entry of B^* . Now assume that the following stronger condition holds for the matrices A and B :

There exist some row matrices c_{ij} , $i, j = 1, \dots, n$ such that each b_{ij} is the sum of the entries of $c_{i,j}$ and each entry of each A_{ij} is a sum of certain entries of c_{ij} such that each entry of c_{ij} appears exactly once as a summand in each row of A_{ij} .

Then the commutative identity implies $A^*\rho = \rho B^*$. By adding 0's to the row matrices c_{ij} we can make all of them size $1 \times k$, for some k , or alternatively, as we do below, we can make each c_{ij} size $1 \times k_i$, so that the size of c_{ij} only depends on i . Thus the row matrices c_{ij} can be arranged in the form of a block matrix $C = (c_{ij})_{ij}$ as below.

Before formally defining the commutative identity, we introduce some notation. Let S be any semiring and consider matrices $A \in S^{m \times n}$ and $B_1, \dots, B_m \in S^{n \times p}$. We let $A\|(B_1, \dots, B_m)$ denote the matrix in $S^{m \times p}$ whose rows are $A_1B_1, \dots, A_mB_m$, where $A_1, \dots, A_m$ are the rows of A .

DEFINITION 5.1 *Suppose that S is a partial $*$ -semiring. We say that the commutative identity holds in S if for all $C \in D(S)^{n \times k}$, $m \times n$ functional matrix ρ , $k \times m$ functional matrices $\rho_1, \dots, \rho_m$ and $k \times n$ functional matrices $\tau_1, \dots, \tau_n$ with $\rho_i\rho = \tau_i\rho$ for all $i = 1, \dots, m$,*

$$((\rho C)\|(\rho_1, \dots, \rho_m))^*\rho = \rho(C\|(\tau_1, \dots, \tau_n))^*.$$

Note that under the assumptions we have $A\rho = \rho B$ for the matrices $A = (\rho C)\|(\rho_1, \dots, \rho_m)$ and $B = C\|(\tau_1, \dots, \tau_n)$, and that the commutative identity asserts that $A^*\rho = \rho B^*$.

The commutative identity has a dual which also holds in all (partial) iteration semirings, see [18, 11]. It can be formulated as follows.

DEFINITION 5.2 *Suppose that S is a partial $*$ -semiring. We say that the dual commutative identity holds in S if for all $C \in D(S)^{k \times n}$, $m \times n$ functional matrix ρ , $k \times m$ functional matrices $\rho_1, \dots, \rho_m$ and $k \times n$ functional matrices $\tau_1, \dots, \tau_n$ with $\rho_i\rho = \tau_i\rho$ for all $i = 1, \dots, m$,*

$$\rho^T((\rho_1^T, \dots, \rho_m^T)\|(C\rho^T))^* = ((\tau_1^T, \dots, \tau_n^T)\|C)^*\rho^T.$$

Here $(B_1, \dots, B_n)\|A$ is the matrix whose columns are $B_1A_1, \dots, B_nA_n$, where $A_1, \dots, A_n$ are the columns of A , A is $m \times n$ and $B_1, \dots, B_n$ are $p \times m$.

DEFINITION 5.3 A semiring S is atomistic if for any $a_1, \dots, a_m$ and $b_1, \dots, b_n$ in S if $a_1 + \dots + a_m = b_1 + \dots + b_n$ then there exist $c_1, \dots, c_k$ in S and partitions $I_1, \dots, I_m$ and $J_1, \dots, J_n$ of the set $\{1, \dots, k\}$ such that

$$\begin{aligned} a_i &= \sum_{p \in I_i} c_p \\ b_j &= \sum_{q \in J_j} c_q \end{aligned}$$

for each $i = 1, \dots, m$ and $j = 1, \dots, n$.

Examples of atomistic semirings are $\mathbb{B}$, $\mathbb{N}$ and $\mathbb{N}_\infty$.

PROPOSITION 5.4 Suppose that S is atomistic and $A \in S^{m \times m}$, $B \in S^{n \times n}$ are such that $A\rho = \rho B$ holds for some $m \times n$ functional matrix ρ . Then there is a matrix $C \in S^{n \times k}$, $k \times m$ functional matrices $\rho_1, \dots, \rho_m$ and $k \times n$ functional matrices $\tau_1, \dots, \tau_n$ with $\rho_i \rho = \tau_i \rho$ for all $i = 1, \dots, m$ such that

$$\begin{aligned} A &= (\rho C) \| (\rho_1, \dots, \rho_m) \\ B &= C \| (\tau_1, \dots, \tau_n). \end{aligned}$$

Proof. It suffices to consider the case when ρ is surjective and monotone. Thus, the assumption is that A is a block matrix $(A_{ij})_{ij}$ such that the sum of each row of each A_{ij} is b_{ij} , the (i, j) th entry of B . Since S is atomistic, for each (i, j) there is a row matrix C_{ij} such that the sum of its entries is b_{ij} and each entry of each row of each A_{ij} can be written as a sum of certain entries of C_{ij} in such a way that each entry of C_{ij} appears exactly once as a summand in each row of A_{ij} . But this is clear since the semiring is atomistic. $\square$

In a similar way, we have:

PROPOSITION 5.5 Suppose that S is atomistic and $A \in S^{m \times m}$, $B \in S^{n \times n}$ are matrices such that $\rho^T A = B \rho^T$ holds for some $m \times n$ functional matrix ρ . Then there is a matrix $C \in S^{k \times n}$, $k \times m$ functional matrices $\rho_1, \dots, \rho_m$ and $k \times n$ functional matrices $\tau_1, \dots, \tau_n$ with $\rho_i \rho = \tau_i \rho$ for all $i = 1, \dots, m$ such that

$$\begin{aligned} A &= (\rho_1^T, \dots, \rho_m^T) \| (C \rho^T) \\ B &= (\tau_1^T, \dots, \tau_n^T) \| C. \end{aligned}$$

PROPOSITION 5.6 Suppose that S is atomistic. Let $A \in S\langle \Sigma \rangle^{m \times m}$, $B \in S\langle \Sigma \rangle^{n \times n}$ be matrices and let ρ be a functional matrix of size $m \times n$. If $A\rho = \rho B$ then there is a matrix $C \in S\langle \Sigma \rangle^{n \times k}$, $k \times m$ functional matrices $\rho_1, \dots, \rho_m$ and $k \times n$ functional matrices $\tau_1, \dots, \tau_n$ with $\rho_i \rho = \tau_i \rho$ for all $i = 1, \dots, m$ such that

$$\begin{aligned} A &= (\rho C) \| (\rho_1, \dots, \rho_m) \\ B &= C \| (\tau_1, \dots, \tau_n). \end{aligned}$$

Proof. There exists a finite $\Sigma_0 \subseteq \Sigma$ such that whenever A or B has an entry which has a summand $s\sigma$ where $\sigma \in \Sigma$ and s is not 0, then $\sigma \in \Sigma_0$. Now for each $\sigma \in \Sigma_0$, let A^σ denote the $m \times m$ matrix whose (i, j) th entry for $i, j = 1, \dots, m$ is $s\sigma$ where s is the coefficient of σ in A_{ij} , the (i, j) th entry of A . If there is no such summand, let $A_{ij}^\sigma = 0$. Define the $n \times n$ matrices B^σ , $\sigma \in \Sigma_0$ in the same way. We then have $A^\sigma \rho = \rho B^\sigma$, for each $\sigma \in \Sigma_0$. Thus, by

Proposition 5.4, for each $\sigma \in \Sigma_0$ there is a matrix $C^\sigma \in S\langle\{\sigma\}\rangle^{m \times k^\sigma}$ and functional matrices ρ_i^σ and τ_j^σ , $i = 1, \dots, m$, $j = 1, \dots, n$ of appropriate size with $\rho_i^\sigma \rho = \tau_{i\rho}^\sigma$ such that

$$\begin{aligned} A^\sigma &= (\rho C^\sigma) \|(\rho_1^\sigma, \dots, \rho_m^\sigma) \\ B^\sigma &= C^\sigma \|(\tau_1^\sigma, \dots, \tau_n^\sigma). \end{aligned}$$

Let $\Sigma_0 = \{\sigma_1, \dots, \sigma_p\}$, say. Define

$$C = \begin{pmatrix} C^{\sigma_1} & \dots & C^{\sigma_p} \end{pmatrix}$$

and

$$\begin{aligned} \rho_i &= \begin{pmatrix} \rho_i^{\sigma_1} \\ \vdots \\ \rho_i^{\sigma_p} \end{pmatrix} \\ \tau_j &= \begin{pmatrix} \tau_j^{\sigma_1} \\ \vdots \\ \tau_j^{\sigma_p} \end{pmatrix} \end{aligned}$$

for all $i = 1, \dots, m$ and $j = 1, \dots, n$. Then for each i ,

$$\begin{aligned} \rho_i \rho &= \begin{pmatrix} \rho_i^{\sigma_1} \\ \vdots \\ \rho_i^{\sigma_p} \end{pmatrix} \rho \\ &= \begin{pmatrix} \rho_i^{\sigma_1} \rho \\ \vdots \\ \rho_i^{\sigma_p} \rho \end{pmatrix} \\ &= \begin{pmatrix} \tau_{i\rho}^{\sigma_1} \\ \vdots \\ \tau_{i\rho}^{\sigma_p} \end{pmatrix} \\ &= \tau_{i\rho}. \end{aligned}$$

Also,

$$\begin{aligned} (\rho C) \|(\rho_1, \dots, \rho_m) &= \begin{pmatrix} C_{1\rho}^{\sigma_1} \rho_1^{\sigma_1} + \dots + C_{1\rho}^{\sigma_p} \rho_1^{\sigma_p} \\ \vdots \\ C_{m\rho}^{\sigma_1} \rho_m^{\sigma_1} + \dots + C_{m\rho}^{\sigma_p} \rho_m^{\sigma_p} \end{pmatrix} \\ &= \begin{pmatrix} A_1^{\sigma_1} + \dots + A_1^{\sigma_p} \\ \vdots \\ A_m^{\sigma_1} + \dots + A_m^{\sigma_p} \end{pmatrix} \\ &= A, \end{aligned}$$

where $A_1, \dots, A_m$ denote the rows of A . In a similar way, $C \|(\tau_1, \dots, \tau_n) = B$. $\square$

Symmetrically, we have:

PROPOSITION 5.7 *Suppose that S is atomistic. Let $A \in S\langle\Sigma\rangle^{m \times m}$, $B \in S\langle\Sigma\rangle^{n \times n}$ and let ρ be a functional matrix of size $m \times n$. If $\rho^T A = B \rho^T$ then there is a matrix $C \in S\langle\Sigma\rangle^{k \times n}$, $k \times m$ functional matrices $\rho_1, \dots, \rho_m$ and $k \times n$ functional matrices $\tau_1, \dots, \tau_n$ with $\rho_i \rho = \tau_{i\rho}$ for all $i = 1, \dots, m$ such that*

$$\begin{aligned} A &= (\rho_1^T, \dots, \rho_m^T) \| (C \rho^T) \\ B &= (\tau_1^T, \dots, \tau_n^T) \| C. \end{aligned}$$

6 Free partial iteration semirings

In this section, our aim is to show that for any set Σ , $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the class of *partial* iteration semirings. For this reason, assume that S is a partial iteration semiring and h is a function $\Sigma \rightarrow D(S)$. We can extend h to a semiring morphism $\mathbb{N}\langle\Sigma^*\rangle \rightarrow S$. In particular, h is defined on $\mathbb{N}$ and on $\mathbb{N}\langle\Sigma\rangle$, and in a pointwise manner, on matrices with entries in $\mathbb{N}$ or $\mathbb{N}\langle\Sigma\rangle$.

We want to show that h can be extended to a unique morphism $h^\sharp : \mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle \rightarrow S$ of partial iteration semirings. For this reason, we will consider automata $\mathbf{A} = (\alpha, A, \beta)$ (in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$) where $\alpha \in \mathbb{N}^{1 \times n}$, $\beta \in \mathbb{N}^{n \times 1}$ and $A \in \mathbb{N}\langle\Sigma\rangle^{n \times n}$ for some n . Using the function h , we define the *image of $\mathbf{A}$* as the automaton $\mathbf{A}h$ in S : $\mathbf{A}h = (\alpha h, Ah, \beta h)$. We know from Theorem 4.3 that we are forced to define $h^\sharp$ by $|\mathbf{A}|h = |\mathbf{A}h|$, for all automata $\mathbf{A}$. We also know that if this function is well-defined, then it is a morphism $\mathbb{N}\langle\langle\Sigma^*\rangle\rangle \rightarrow S$ of partial iteration semirings (which clearly extends h). So all we have to show is that $h^\sharp$ is well-defined. The proof of this fact relies on a result proved in [4] that we recall now.

DEFINITION 6.1 *Let $\mathbf{A} = (\alpha, A, \beta)$ and $\mathbf{B} = (\gamma, B, \delta)$ be two automata (in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$) of dimension m and n , respectively. We say that an $m \times n$ functional matrix ρ is a *simulation* $\mathbf{A} \rightarrow \mathbf{B}$ if $\alpha\rho = \gamma$, $\rho\delta = \beta$ and $A\rho = \rho B$ hold. Moreover, we say that ρ is a *dual simulation* $\mathbf{A} \rightarrow \mathbf{B}$ if ρ is a simulation $\mathbf{A}^T \rightarrow \mathbf{B}^T$, where $\mathbf{A}^T = (\beta^T, A^T, \alpha^T)$ and $\mathbf{B}^T$ is defined in the same way.*

Note that ρ is a dual simulation $\mathbf{A} \rightarrow \mathbf{B}$ iff $\gamma\rho^T = \alpha$, $B\rho^T = \rho^T A$ and $\rho^T\beta = \delta$ hold.

(More general simulations were defined in [8]. The simulations defined above are the functional and dual functional simulations of [8]. In the papers [3, 4], the terms “covering” and “co-covering” are used for simulation and dual simulation. Moreover, only simulations and dual simulations corresponding to surjective functions are considered, since in the formulation of Theorem 6.2 given in [4], the automata are “trim”, i.e., without useless states.) Let $\sim$ denote the least equivalence relation such that $\mathbf{A} \sim \mathbf{B}$ holds whenever there is a functional simulation or a dual functional simulation $\mathbf{A} \rightarrow \mathbf{B}$. Moreover, call two automata $\mathbf{A}$ and $\mathbf{B}$ *equivalent* if $|\mathbf{A}| = |\mathbf{B}|$. The following result was proved in [4]:

THEOREM 6.2 *Two automata $\mathbf{A}$ and $\mathbf{B}$ in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ are equivalent iff $\mathbf{A} \sim \mathbf{B}$.*

So our task reduces to showing that for automata $\mathbf{A}$ and $\mathbf{B}$ in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, if there is a functional or a dual functional simulation $\mathbf{A} \rightarrow \mathbf{B}$, then $|\mathbf{A}h| = |\mathbf{B}h|$.

LEMMA 6.3 *Suppose that $\mathbf{A} = (\alpha, A, \beta)$ and $\mathbf{B} = (\gamma, B, \delta)$ are automata in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ of dimension m and n , respectively. Suppose that ρ is an $m \times n$ functional matrix which is a simulation $\mathbf{A} \rightarrow \mathbf{B}$. Then $|\mathbf{A}h| = |\mathbf{B}h|$.*

Proof. Since $A\rho = \rho B$, it follows from Proposition 5.6 that there exists a matrix $C \in \mathbb{N}\langle\Sigma\rangle^{n \times k}$ and $k \times m$ functional matrices $\rho_1, \dots, \rho_m$ and $k \times n$ functional matrices $\tau_1, \dots, \tau_n$ with $\rho_i\rho = \tau_i\rho$ for all i such that $A = (\rho C)\|(\rho_1, \dots, \rho_m)$ and $B = C\|(\tau_1, \dots, \tau_n)$. Thus, also $Ah = ((\rho h)(Ch))\|(\rho_1 h, \dots, \rho_m h)$ and $Bh = (Ch)\|(\tau_1 h, \dots, \tau_n h)$. Thus, by the commutative iden-

tity, $(Ah)^*(\rho h) = (\rho h)(Bh)^*$. Thus,

$$\begin{aligned}
|\mathbf{A}h| &= (\alpha h)(Ah)^*(\beta h) \\
&= (\alpha h)(Ah)^*((\rho\delta)h) \\
&= (\alpha h)(Ah)^*(\rho h)(\delta h) \\
&= (\alpha h)(\rho h)(Bh)^*(\delta h) \\
&= ((\alpha\rho)h)(Bh)^*(\delta h) \\
&= (\gamma h)(Bh)^*(\delta h) \\
&= |\mathbf{B}h|. \quad \square
\end{aligned}$$

LEMMA 6.4 *Suppose that $\mathbf{A}$ and $\mathbf{B}$ are finite automata as above of dimension m and n , respectively. Suppose that ρ is an $m \times n$ functional matrix which is a dual simulation $\mathbf{A} \rightarrow \mathbf{B}$. Then $|\mathbf{A}h| = |\mathbf{B}h|$.*

Proof. Since ρ is a simulation $\mathbf{A}^T \rightarrow \mathbf{B}^T$, it follows as above that $A^T = (\rho C^T) \|(\rho_1, \dots, \rho_m)$ and $B^T = C^T \|(\tau_1, \dots, \tau_n)$ for some C^T , $\rho_1, \dots, \rho_m$ and $\tau_1, \dots, \tau_n$ with $\rho_i \rho = \tau_i \rho$. Thus, $A = (\rho_1^T, \dots, \rho_m^T) \| (C \rho^T)$ and $B = (\tau_1^T, \dots, \tau_n^T) \| C$. The proof can be completed as above using the dual commutative identity. $\square$

The main result of this section is:

THEOREM 6.5 *$\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the class of partial iteration semirings. In detail, given any partial iteration semiring S and function $h : \Sigma \rightarrow D(S)$, there is a unique partial iteration semiring morphism $h^\sharp : \mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle \rightarrow S$ extending h .*

Proof. Given S and h , define $h^\sharp$ as follows. First, extend h to a semiring morphism $\mathbb{N}\langle\Sigma^*\rangle \rightarrow S'$. By Theorem 4.2, we know that every rational series in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is the behavior of an automaton in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$. We also know that for any rational power series $r \in \mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ recognized by an automaton $\mathbf{A}$, we are forced to define $rh^\sharp = |\mathbf{A}h|$. By Theorem 6.2, Lemma 6.3 and Lemma 6.4, $h^\sharp$ is well-defined. It is clear that $h^\sharp$ extends h . Moreover, by Theorem 4.3, $h^\sharp$ is a morphism of partial iteration semirings. $\square$

REMARK 6.6 The paper [11] also defines *partial iterative semirings* as partial $*$ -semirings S such that for each $a, b \in S$, if $a \in D(S)$, then a^*b is the *unique* solution of the equation $x = ax + b$. It is shown that every partial iterative semiring is a partial iteration semiring, and that for any semiring S and set Σ , the power series semiring $S\langle\langle\Sigma^*\rangle\rangle$ is a partial iterative semiring. Thus, $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is also a partial iterative semiring. Since morphisms of partial iterative semirings preserve star, it follows that $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is the free partial iterative semiring on Σ . This fact is related to a result proved in [29], where Morisaki and Sakai extended Salomaa's axiomatization [32] of regular languages to rational power series over *fields* (or more generally, *principal ideal domains*).

Theorem 6.5 can be generalized. Consider a power series semiring $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ where S is any semiring. We can define simulations and dual simulations and the relation $\sim$ for automata in $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ over (S, Σ) in the same way as above. For example, when $\mathbf{A} = (\alpha, A, \beta)$ and $\mathbf{B} = (\gamma, B, \delta)$ are automata over (S, Σ) of dimension m and n , then a simulation $\mathbf{A} \rightarrow \mathbf{B}$ is an $m \times n$ functional matrix ρ such that $\alpha\rho = \gamma$, $A\rho = \rho B$ and $\beta = \rho\delta$. If ρ is a simulation $\mathbf{A} \rightarrow \mathbf{B}$, then

$$\alpha A^k \beta = \alpha A^k \rho \delta = \alpha \rho B^k \delta = \gamma B^k \delta$$

for all k , and thus $|\mathbf{A}| = |\mathbf{B}|$, i.e., $\mathbf{A}$ and $\mathbf{B}$ are equivalent. In a similar way, if ρ is a dual simulation $\mathbf{A} \rightarrow \mathbf{B}$, then $|\mathbf{A}| = |\mathbf{B}|$. Thus, if $\mathbf{A} \sim \mathbf{B}$, then $\mathbf{A}$ and $\mathbf{B}$ are equivalent. In the following generalization of Theorem 6.5 we will assume that also the converse property is true, if $\mathbf{A}$ and $\mathbf{B}$ are equivalent then $\mathbf{A} \sim \mathbf{B}$.

THEOREM 6.7 *Let S be a semiring and Σ a set. Suppose that if two automata $\mathbf{A}, \mathbf{B}$ in $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ over (S, Σ) are equivalent then $\mathbf{A} \sim \mathbf{B}$ holds. Moreover, suppose that S is atomistic. Then $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ has the following universal property. Given any partial iteration semiring S' , semiring morphism $h_S : S \rightarrow S'$ and function $h : \Sigma \rightarrow S'$ such that sh_S commutes with ah for all $s \in S$ and $a \in \Sigma$, there is a unique partial iteration semiring morphism $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle \rightarrow S'$ extending h_S and h .*

The proof is exactly the same. Theorem 6.7 is applicable for example to the boolean semiring $\mathbb{B}$ (see [8]), and the semirings $\mathbf{k}$ defined in Section 7. However, for rings simpler characterizations exist, cf. [5].

REMARK 6.8 Without the assumption that S is atomistic, we only have the following fact. Suppose that S' is a partial Conway semiring satisfying the *functorial star implications* [8]

$$\begin{aligned} A\rho = \rho a &\Rightarrow A^*\rho = \rho a^* \\ \rho^T A = a\rho^T &\Rightarrow \rho^T A^* = a^*\rho^T \end{aligned}$$

for all $a \in D(S')$ and $A \in S'^{m \times m}$ whose entries are in $D(S')$, and for all $m \times 1$ functional matrices, $m \geq 2$. Then, as shown in [8, 11], S' is a partial iteration semiring satisfying the functorial star implications

$$\begin{aligned} A\rho = \rho B &\Rightarrow A^*\rho = \rho B^* \\ \rho^T A = B\rho^T &\Rightarrow \rho^T A^* = B^*\rho^T \end{aligned}$$

for all $A \in S'^{m \times m}$, $B \in S'^{n \times n}$ whose entries are in $D(S')$, and for all $m \times n$ functional matrices ρ for any integers m, n . As above, suppose that if two automata $\mathbf{A}, \mathbf{B}$ in $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ over (S, Σ) are equivalent then $\mathbf{A} \sim \mathbf{B}$. Then for any semiring morphism $h_S : S \rightarrow S'$ and function $h : \Sigma \rightarrow S'$ such that $\Sigma h \subseteq D(S')$ and sh_S commutes with ah , for all $s \in S$ and $a \in \Sigma$, there is a unique partial iteration semiring morphism $S^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle \rightarrow S'$ extending h_S and h .

7 A characterization

We have seen that for any set Σ , $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the class of *partial* iteration semirings. In particular, $\mathbb{N}$ is initial in the class of partial iteration semirings. This latter fact is also clear by noting that the star operation is completely undefined in $\mathbb{N}$ and that $\mathbb{N}$ is initial in the class of semirings. The smallest iteration semiring which contains $\mathbb{N}$ as a subsemiring is $\mathbb{N}_\infty$, the completion of $\mathbb{N}$ with a point of infinity denoted ∞ and star operation defined by $0^* = 1$ and $n^* = \infty$ for all $n \neq 0$. In this section our aim is to show that the iteration semirings $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ are the free algebras in a subvariety of iteration semirings defined by a few simple identities. By Proposition 3.8, $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is an iteration semiring.

The structure of the initial iteration semiring was described in [8]. Its elements are

$$0, 1, 2, \dots, 1^*, (1^*)^2, \dots, 1^{**},$$

ordered as indicated. Sum and product on the integers are the standard operations; the sum and product on the remaining elements are given by:

$$\begin{aligned} x + y &= \max\{x, y\}, \quad \text{if } x \geq 1^* \text{ or } y \geq 1^* \\ (1^*)^n (1^*)^p &= (1^*)^{n+p} \\ x 1^{**} = 1^{**} x &= 1^{**}, \quad \text{if } x \neq 0. \end{aligned}$$

Lastly, the star operation is defined by:

$$x^* = \begin{cases} 1 & \text{if } x = 0 \\ 1^* & \text{if } x = 1 \\ 1^{**} & \text{otherwise.} \end{cases}$$

Identifying 1^* and 1^{**} , the resulting congruence collapses the elements

$$1^*, (1^*)^2, \dots, 1^{**},$$

so that the corresponding quotient $*$ -semiring is isomorphic to $\mathbb{N}_\infty$.

In this section we will characterize the iteration semirings $\mathbb{N}_\infty^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$ as the free algebras in the subvariety $\mathcal{V}$ of iteration semirings specified by the following identities:

$$1^* 1^* = 1^* \tag{11}$$

$$1^* a = a 1^* \tag{12}$$

$$1^* (1^* a)^* = 1^* a^*. \tag{13}$$

PROPOSITION 7.1 *The identity $1^* = 1^{**}$ holds in $\mathcal{V}$.*

Proof. Instantiating (13) with $a = 1$ and using (11) we have $1^* 1^{**} = 1^* 1^* = 1^*$. But by the above description of the initial iteration semiring, $1^* 1^{**} = 1^{**}$ in any iteration semiring. $\square$

Since $2^* = 3^* = \dots = 1^{**}$ in the initial iteration semiring, it follows that $n^* = 1^*$ holds in $\mathcal{V}$ for any integer $n \geq 1$ viewed as a term.

Thus, (11) may be replaced by the identity $1^* = 1^{**}$. Also, by (8), $(1 + a)^* = (1^* a)^* 1^*$, so that in view of (12), equation (13) is equivalent to

$$(1 + a)^* = 1^* a^*. \tag{14}$$

More generally, we have that

$$(n + a)^* = 1^* a^* \tag{15}$$

holds in $\mathcal{V}$, for any $n \in \mathbb{N}_\infty$, $n \neq 0$ viewed as a term. Also, in view of the other axioms, (13) is equivalent to the simpler

$$a^{**} = 1^* a^* \tag{16}$$

since $(a + 1)^* = 1^* (a 1^*)^* = 1^* + 1^* (a 1^*)^+ = 1 + 1^* + 1^* (a 1^*)^+ = 1 + (a + 1)^* = a^* a^{**} + 1 = a^{**}$ using only the Conway identities.

Since $\mathbb{N}_\infty$ satisfies the identities (11), (12) and (13), and since $1^* = 1^{**}$ holds in $\mathcal{V}$, we have:

COROLLARY 7.2 *$\mathbb{N}_\infty$ is initial in $\mathcal{V}$.*

Also, for each set Σ , both $\mathbb{N}_\infty \langle\langle \Sigma^* \rangle\rangle$ and $\mathbb{N}_\infty^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$ are in $\mathcal{V}$.

Consider the set of *iteration semiring terms*, or just *terms* over Σ defined by

$$t = 0 \mid 1 \mid a, a \in \Sigma \mid t + t \mid t \cdot t \mid t^*.$$

A term is called *constant term* or just *constant* if it contains no occurrence of any letter in Σ . We will say that two terms s, t are *equivalent* if they are equivalent modulo the defining identities of $\mathcal{V}$, i.e., when the identity $s = t$ holds in $\mathcal{V}$.

Each term t over Σ evaluates to a series $|t|$ in $\mathbb{N}_\infty^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$ as usual. Since $\mathbb{N}_\infty$ is initial in $\mathcal{V}$, for any constant terms s, t we have $|s| = |t|$ iff $s = t$ holds in $\mathcal{V}$. We may thus identify each constant term with an element of $\mathbb{N}_\infty$.

The class I of *ideal terms* is the least class of terms with the following properties.

1. $0 \in I$ and $a \in I$ for all $a \in \Sigma$.
2. If $s \in I$ and $t \in I$ then $s + t \in I$.
3. If $s \in I$ and $t \in I$ or t is a constant in $\mathbb{N}$, then st and ts are in I .
4. If $s \in I$ then s^+ is in I , where s^+ is an abbreviation for ss^* .

LEMMA 7.3 *When t is ideal, $|t|$ is proper and $|t| \in \mathbb{N} \langle\langle \Sigma^* \rangle\rangle$.*

The easy proof is omitted. It then follows that each ideal term t also evaluates to a series in the partial iteration semiring $\mathbb{N}_\infty^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$, and that this series is the same as the evaluation of t in $\mathbb{N}_\infty^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$.

LEMMA 7.4 *For every term t there is an equivalent term of the form $t_c + t_0 + 1^*t_\infty$, where t_c is a constant in $\mathbb{N}$, t_0 is an ideal term, and t_∞ is a term. Moreover, if $t_c \neq 0$ then t_∞ is ideal.*

Proof. This fact is implied by the following claim:

For every term t there is an equivalent term of the form $t_c + t_0 + 1^*t_\infty$, where t_c is a constant in $\mathbb{N}_\infty$, t_0 and t_∞ are ideal terms.

We prove this fact by induction on the structure of t . When t is $0, 1$ or a letter in Σ , our claim is clear. Suppose that $t = p + s$. Then t is equivalent to $(p_c + s_c) + (p_0 + s_0) + 1^*(p_\infty + s_\infty)$. Assume now that $t = ps$. Then using (11) and (12), t is equivalent to $p_cs_c + (p_cs_0 + p_0s_c) + 1^*((p_c + p_0)s_\infty + p_\infty(s_c + s_0) + p_\infty s_\infty)$. Finally, assume that $t = s^*$. If $s_c = 0$, then t is equivalent to $1 + s_0^+ + 1^*(s_0 + s_\infty)^* s_\infty s_0^*$ as shown by the following computation using the sum star and product star identities and (11), (12) and (13).

$$\begin{aligned} (s_0 + 1^*s_\infty)^* &= s_0^*(1^*s_\infty s_0^*)^* \\ &= s_0^*(1 + 1^*(s_\infty s_0^* 1^*)^* s_\infty s_0^*) \\ &= s_0^* + s_0^* 1^*(1^*s_\infty s_0^*)^* s_\infty s_0^* \\ &= s_0^* + s_0^* 1^*(s_\infty s_0^*)^* s_\infty s_0^* \\ &= s_0^* + 1^*s_0^*(s_\infty s_0^*)^* s_\infty s_0^* \\ &= 1 + s_0^+ + 1^*(s_0 + s_\infty)^* s_\infty s_0^*. \end{aligned}$$

If $s_c \neq 0$, then using (15) we have that t is equivalent to $1^*(s_0 + s_\infty)^* = 1^* + 1^*(s_0 + s_\infty)^+$. In either case, s^* is of the required form. $\square$

As an immediate corollary, we note the following Fatou property:

COROLLARY 7.5 *If $s \in \mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ and all coefficients of s are in $\mathbb{N}$, then $s \in \mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$.*

In our proof that each iteration semiring $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the variety $\mathcal{V}$ we will make use of the corresponding fact for the boolean semiring, proved in Krob [25].

THEOREM 7.6 *For each Σ , $\mathbb{B}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the variety of all iteration semirings satisfying $1^* = 1$.*

Let $\mathcal{W}$ denote the variety of iteration semirings satisfying $1^* = 1$. It is clear that $\mathcal{W}$ is a subvariety of $\mathcal{V}$. We introduce a construction which assigns to every iteration semiring A in $\mathcal{V}$ an iteration semiring 1^*A in $\mathcal{W}$.

Suppose that $A \in \mathcal{V}$. We define $1^*A = \{1^*a : a \in A\}$. It is clear that 1^*A contains 0 and is closed under sum and product. Also, using (11), (12) and (13), $(1^*a)^+ = 1^*a(1^*a)^* = 1^*a1^*a^* = 1^*a^+$, showing that 1^*A is closed under the “plus operation” $a \mapsto a^+$. However, 1^*A does not necessarily contain 1 and is not necessarily closed under star.

DEFINITION 7.7 *For each $A \in \mathcal{V}$, we equip 1^*A with the following operations and constants. The sum $+$ and product $\cdot$ operations and the constant 0 are inherited from A , the constant $\mathbf{1}$ is 1^* and the star operation $^\otimes$ is defined by $(1^*a)^\otimes = 1^*(1^*a)^*$.*

Note that $^\otimes$ is well-defined, since if $1^*a = 1^*b$, for some $a, b \in A$, then $1^*(1^*a)^* = 1^*(1^*b)^*$. Also, by (13), $(1^*a)^\otimes = 1^*a^*$. Using this, it follows that the plus operation of 1^*A determined by the star operation $^\otimes$ is the restriction of the plus operation of A . Indeed, for all $a \in A$, $1^*a(1^*a)^\otimes = 1^*a1^*a^* = 1^*a^+ = (1^*a)^+$.

LEMMA 7.8 *For any $A \in \mathcal{V}$, the assignment $h : a \mapsto 1^*a$, $a \in A$ preserves all operations and constants.*

Proof. Clearly, we have $0h = 0$ and $1h = \mathbf{1}$. Also, $(a+b)h = 1^*(a+b) = 1^*a + 1^*b = ah + bh$ and $(ab)h = 1^*ab = 1^*1^*ab = 1^*a1^*b = (ah)(bh)$, for all $a, b \in A$. Finally, $a^*h = 1^*a^* = (1^*a)^\otimes = (ah)^\otimes$. $\square$

COROLLARY 7.9 *For each A in $\mathcal{V}$, 1^*A is an iteration semiring in $\mathcal{W}$.*

Proof. Since the morphism h is surjective, we have $1^*A \in \mathcal{V}$. Since also $\mathbf{1}^\otimes = 1^*1^{**} = 1^*1^* = 1^* = \mathbf{1}$, it holds that $1^*A \in \mathcal{W}$. $\square$

For the next corollary, note that if t is a term over Σ and A is an iteration semiring, then t induces a function $A^\Sigma \rightarrow A$ as usual. We will denote this function by t^A . Below we will write function composition in the diagrammatic order.

COROLLARY 7.10 *Suppose that t is a term over Σ and $A \in \mathcal{V}$. Then $(1^*t)^A$ can be factored as $\bar{h} \circ (1^*t)^{1^*A}$, where h is the morphism $A \rightarrow 1^*A$ of Lemma 7.8 and $\bar{h} : A^\Sigma \rightarrow (1^*A)^\Sigma$, $e \mapsto e \circ h$.*

Proof. Since h is a morphism,

$$\bar{h} \circ (1^*t)^{1^*A} = (1^*t)^A \circ h = 1^*(1^*t)^A = (1^*1^*)t^A = (1^*t)^A. \quad \square$$

COROLLARY 7.11 *Suppose that s, t are terms over Σ and $A \in \mathcal{V}$. Then $1^*t = 1^*s$ holds in A iff it holds in 1^*A .*

We will use Theorem 7.6 in the following way. Let Σ be a set and consider a term t over Σ . It is easy to see by induction that if 1^*t evaluates to a series r in $\mathbb{B}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, then in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ it evaluates to the series whose nonzero coefficients are all ∞ and whose support is the same as that of r .

We claim that if 1^*t and 1^*s evaluate to the same series in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, then the identity $1^*t = 1^*s$ holds in $\mathcal{V}$. Let $\mathcal{W}$ denote the variety of iteration semirings satisfying $1^* = 1$. Since 1^*t and 1^*s evaluate to the same series in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, they evaluate to the same series in $\mathbb{B}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$. Thus, by Theorem 7.6, $1^*t = 1^*s$ holds in $\mathcal{W}$. Let $A \in \mathcal{V}$. By Corollary 7.9, $1^*A \in \mathcal{W}$, so $1^*t = 1^*s$ holds in 1^*A . By Corollary 7.11, this implies that $1^*t = 1^*s$ holds in A . Since A was an arbitrary iteration semiring in $\mathcal{V}$, this means that $1^*t = 1^*s$ holds in $\mathcal{V}$.

LEMMA 7.12 *Suppose that t, s are terms over Σ such that the support of $|t|$ is included in the support of $|1^*s|$. Then $t + 1^*s$ is equivalent to 1^*s .*

Proof. By the above argument, $1^*s = 1^*(t + s) = 1^*t + 1^*s$ holds in $\mathcal{V}$. Also, $t + 1^*t = (1 + 1^*)t = 1^*t$ holds. Thus,

$$t + 1^*s = t + 1^*t + 1^*s = 1^*t + 1^*s = 1^*s$$

holds. □

We now prove a stronger version of Lemma 7.4.

LEMMA 7.13 *For every term t there is an equivalent term of the form $t_c + t_0 + 1^*t_\infty$, where t_c is a constant in $\mathbb{N}$, t_0 is an ideal term, and t_∞ is a term. Moreover, if $t_c \neq 0$ then t_∞ is ideal and $|t_0|$ and $|1^*t_\infty|$ have disjoint supports.*

Proof. We know from Lemma 7.4 that t is equivalent to a term of the form $t_c + t_0 + 1^*t_\infty$, where $t_c \in \mathbb{N}$, t_0 is an ideal term and if t_c is not 0 then t_∞ is also ideal. Now $\text{supp}(1^*t_\infty) = \text{supp}(t_\infty)$ is a regular language which we denote by R . Consider the rational series $s_0 = |t_0|$ and write it as the sum $s_1 + s_2$, where $(s_1, w) = (s_0, w)$ if $w \notin R$ and $(s_1, w) = 0$ otherwise, moreover, $(s_2, w) = (s_0, w)$ if $w \in R$ and $(s_2, w) = 0$ otherwise. It is known that s_1 and s_2 are rational (see [5]) and thus there exist ideal terms t_1 and t_2 with $|t_1| = s_1$ and $|t_2| = s_2$. Since $|t_1 + t_2| = s_1 + s_2 = s = |t_0|$, and since these terms are ideal, by Theorem 6.5 we have that $t_0 = t_1 + t_2$ holds in $\mathcal{V}$. Since the support of $|t_2|$ is included in the support of $|1^*t_\infty|$, $t_2 + 1^*t_\infty = 1^*t_\infty$ also holds in $\mathcal{V}$. Summing up, t is equivalent to $t_c + t_0 + 1^*t_\infty$ which is in turn equivalent to $t_c + t_1 + 1^*t_\infty$ proving the claim. □

THEOREM 7.14 *For each set Σ , $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in $\mathcal{V}$.*

Proof. We have already noted that $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is in $\mathcal{V}$. By definition, Σ generates $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$. But we still have to show that if two terms over Σ evaluate to the same series, then they are equivalent. But any term t is equivalent to some term of the form $t_c + t_0 + 1^*t_\infty$ where t_c is a constant in $\mathbb{N}$ and t_0 is ideal, and if $t_c > 0$, then t_∞ is ideal. Now $|t| = |t_c| + |t_0| + |1^*t_\infty|$, where $|t_c| \in \mathbb{N}$, $|t_0| \in \mathbb{N}\langle\langle\Sigma^*\rangle\rangle$ and $|1^*t_\infty| \in \{0, \infty\}\langle\langle\Sigma^*\rangle\rangle$, i.e., each coefficient of the series $|1^*t_\infty|$ is 0 or ∞ . Moreover, $|t_0|$ and $|1^*t_\infty|$ have disjoint supports, and either $|t_c| = 0$ or $|1^*t_\infty|$ is proper. Thus, if $|t| = |s|$, then $|t_c| = |s_c|$, $|t_0| = |s_0|$ and $|1^*t_\infty| = |1^*s_\infty|$. By Corollary 7.2 we have that $t_c = s_c$ holds in $\mathcal{V}$. Since t_0 and s_0 evaluate to the same series in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, by Theorem 6.5 we have that $t_0 = s_0$ holds in $\mathcal{V}$. Finally, by the above discussion, $1^*t_\infty = 1^*s_\infty$ holds in $\mathcal{V}$, proving that $t = s$ holds. □

COROLLARY 7.15 *A series $s \in \mathbb{N}_\infty \langle\langle \Sigma^* \rangle\rangle$ is in $\mathbb{N}_\infty^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$ iff $s = s_0 + s_\infty$ where $s_0 \in \mathbb{N}^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$ and all nonzero coefficients of $s_\infty \in \mathbb{N}_\infty^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$ are equal to ∞ . The series s_0 and s_∞ may be chosen so that they have disjoint supports. Moreover, a series s , all of whose nonzero coefficients are equal to ∞ , is rational iff its support is regular.*

REMARK 7.16 The variety $\mathcal{V}$ is not finitely based, since it has a non-finitely based subvariety $\mathcal{W}$ which has a finite relative axiomatization over $\mathcal{V}$ by the single identity $1^* = 1$. See also [26]. Likewise, the variety of all iteration semirings is non-finitely based.

Recall from [16] that a *complete semiring* is a semiring S which is equipped with a summation operation $\sum_{i \in I} s_i$ for all index sets I satisfying $\sum_{i \in \emptyset} s_i = 0$, $\sum_{i \in \{1,2\}} s_i = s_1 + s_2$, moreover, product distributes over all sums and summation is associative:

$$\begin{aligned} a(\sum_{i \in I} b_i) &= \sum_{i \in I} ab_i \\ (\sum_{i \in I} b_i)a &= \sum_{i \in I} b_i a \\ \sum_{j \in J} \sum_{i \in I_j} a_i &= \sum_{i \in \cup_{j \in J} I_j} a_i, \end{aligned}$$

where in the last equation the sets I_j are pairwise disjoint. *Countably complete semirings* are defined in the same with the additional constraint that all sums are at most countable. Clearly, every complete semiring is countably complete.

An ω -continuous semiring [8] is a semiring S equipped with a partial order such that S is an ω -complete partial order (ω -cpo) with bottom element 0 and the sum and product operations are continuous, i.e., they preserve the suprema of ω -chains. A *continuous semiring* is defined in the same way, it is a cpo with continuous operations such that 0 is the bottom element. Each ω -continuous semiring is a countably complete semiring with

$$\sum_{i \in I} s_i = \sup \{ \sum_{i \in F} s_i : F \subseteq I \text{ finite} \}.$$

Similarly, each continuous semiring is complete. The semiring $\mathbb{N}_\infty$, equipped with the natural order, is continuous. It is well-known that equipped with the pointwise order, $\mathbb{N}_\infty \langle\langle \Sigma^* \rangle\rangle$ is also continuous for each Σ .

When S is countably complete, we can define a star operation on S by $a^* = \sum_{n \geq 0} a^n$. Since ω -continuous, continuous and complete semirings are all countably complete, the same definition applies to these semirings. We point out that the $*$ -semirings so obtained are all in $\mathcal{V}$. Indeed, it is known that when S is countably complete, then S is an iteration semiring (cf. [8]). We have that 1^* is a countable sum of 1 with itself. Using distributivity, it follows that $1^*1^* = 1^*$. By distributivity, we also have (12). Finally, 1^*a^* and $1^*(1^*a)^*$ are both equal to a countable sum $\sum_{i \in I} s_i$ containing for each n a countable number of summands s_i equal to a^n .

By the above observations and the fact that the semirings $\mathbb{N}_\infty \langle\langle \Sigma^* \rangle\rangle$ are continuous and contain the semirings $\mathbb{N}_\infty^{\text{rat}} \langle\langle \Sigma^* \rangle\rangle$, we immediately have:

COROLLARY 7.17 *Continuous, ω -continuous, complete and countably complete semirings, equipped with the above star operation, satisfy exactly the identities of the variety $\mathcal{V}$.*

REMARK 7.18 The set $\mathbb{N}_\infty$ carries another important semiring structure. Equipped with minimum as addition and addition as multiplication (and ∞ as the additive identity element and 0

as the multiplicative identity), $\mathbb{N}_\infty$ is called the *tropical semiring*. It is known that the tropical semiring has a non-finitely based equational theory, cf. [1]. Krob [27] has shown that the equality problem for rational power series in two or more letters over the tropical semiring is undecidable. Rational power series in a single letter over the tropical semiring were treated in [14].

We end this section by pointing out how Theorem 7.6 can be derived from Theorem 7.14. When $k \geq 1$ is an integer, let $\mathbf{k}$ denote the quotient of the iteration semiring $\mathbb{N}_\infty$ obtained by collapsing k and ∞ and thus all elements of $\mathbb{N}_\infty$ at least k . When $k = 1$, $\mathbf{k}$ is just the Boolean semiring $\mathbb{B}$ with star operation $0^* = 1^* = 1$. Our result is:

THEOREM 7.19 *For each integer $k \geq 1$, $\mathbf{k}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the variety of iteration semiring satisfying the identity $1^* = k$.*

Of course, in the statement of the Theorem, k also denotes the term $1 + \dots + 1$ (k times). Since any iteration semiring satisfying $1^* = k$ satisfies (11), (12) and (13), Theorem 7.19 is immediate from Theorem 7.14 if we can show that

the least congruence $\sim$ on $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ which collapses k and 1^*

collapses any rational series in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ with a series all of whose coefficients are either less than k or equal to $1^*(= \infty)$. By Corollary 7.15, it is sufficient to prove this for rational series in $\mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$. The rest of this section is devoted to proving this fact.

LEMMA 7.20 *Suppose that $s \in \mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ such that any nonzero coefficient of s is at least k . Then $s \sim 1^*s$.*

Proof. Let $R = \text{supp}(s)$ which is a regular language in Σ^* (cf. [5]), and let r denote the characteristic series of R , so that for any word w , $(r, w) = 1$ if $w \in R$ and $(r, w) = 0$ otherwise. It is known that r is rational (this is true for any semiring, cf. [5]) and thus kr is also rational. Now it is known that $t = s - kr$ is also rational, see Theorem 1.8 in Chapter VII of [5]. It is clear that $kr \sim 1^*r = 1^*s$. Using this, we have:

$$s = kr + t \sim 1^*s + t = 1^*s. \quad \square$$

PROPOSITION 7.21 *For each integer k and each $s \in \mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ there is a series $r \in \mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ with $s \sim r$ such that all coefficients of r are either less than k or equal to 1^* .*

Proof. In our argument, we will make use of the following known fact from [5]. Given any rational series $s \in \mathbb{N}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, s can be written as a sum of rational series $s_0 + \dots + s_k$ such that each coefficient of any s_i with $i < k$ is 0 or i , and each coefficient of s_k is 0 or $\geq k$. By the previous lemma, $s_k \sim 1^*s_k$, and thus s_k is congruent to the rational series s'_k such that $(s'_k, w) = 1^*$ if $(s_k, w) \geq k$ and $(s'_k, w) = 0$ otherwise. We conclude that $s \sim s_0 + \dots + s_{k-1} + s'_k$ which has the desired property. $\square$

8 A second characterization

In the previous section, we have characterized the semirings $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ as the free algebras in a non-finitely based variety $\mathcal{V}$ of $*$ -semirings. Since $\mathbb{N}_\infty$ has a natural order, $\mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle$ may be

equipped with the pointwise order. This order on $\mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle$ is actually the same as the sum order: For all series $s, s' \in \mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle$, $s \leq s'$ iff there is a series r with $s + r = s'$. Moreover, since $\mathbb{N}_\infty$ is a continuous semiring, cf. e.g. [19], so is $\mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle$. In particular, any map $x \mapsto sx + r$ over $\mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle$ has the series s^*r as its least *pre-fixed point* (since $ss^*r + r \leq s^*r$ and for all s' , if $ss' + r \leq s'$ then $s^*r \leq s'$). Moreover, rs^* is the least pre-fixed point of the map $x \mapsto xs + r$. The semiring $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, equipped with the pointwise order inherited from $\mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle$ also has these least pre-fixed point properties. However, in the main result of this section, we will have to work with the sum order on $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ which is not the same as the pointwise order. It is known that for $r, s \in \mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ with $r \leq s$ in the pointwise order, the difference $s - r$ may *not* be rational (see [5]), so that there may not exist a *rational* series r' with $r + r' = s$. Since $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle = \mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle \cap \mathbb{N}\langle\langle\Sigma^*\rangle\rangle$, the same holds for $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$. But the above least pre-fixed point property still holds in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ with the sum order, as will be shown below.

For the rest of this paper, by an *ordered semiring* we shall mean a semiring S equipped with a partial order $\leq$ preserved by sum and product: If $a \leq a'$ and $b \leq b'$ then $a + b \leq a' + b'$ and $ab \leq a'b'$. Following [19], we call a $*$ -semiring an *inductive $*$ -semiring* if it is an ordered semiring such that the following hold for all $a, b, x \in S$:

$$aa^* + 1 \leq a^* \quad (17)$$

$$ax + b \leq x \quad \Rightarrow \quad a^*b \leq x. \quad (18)$$

It then follows that for any a, b , a^*b is the least pre-fixed point of the map $x \mapsto ax + b$, and is actually a fixed point. Moreover, it is known that the star operation is also monotone in any inductive $*$ -semiring. A *symmetric inductive $*$ -semiring* S also satisfies

$$xa + b \leq x \quad \Rightarrow \quad ba^* \leq x \quad (19)$$

for all $a, b, x \in S$. In [23], Kozen defines a *Kleene algebra* as an idempotent symmetric inductive $*$ -semiring. (Note that if an ordered semiring S is idempotent, then the partial order is the semilattice order: $a \leq b$ iff $a + b = b$.) A morphism of (symmetric) inductive $*$ -semirings is a $*$ -semiring morphism which preserves the order.

The following result was proved in [19]:

THEOREM 8.1 *Every inductive $*$ -semiring is an iteration semiring satisfying $1^* = 1^{**}$.*

We call a (symmetric) inductive $*$ -semiring *sum ordered*, if its order relation is given by $a \leq b$ iff there is some c with $a + c = b$. Suppose that S is an inductive $*$ -semiring. Since for any $x \in S$, $1x + 0 = x$, we have that $0 = 1^*0 \leq x$. Thus, 0 is the least element of S and since the order is preserved by addition, $x \leq x + y$ for all $x, y \in S$, so that the order on S is an extension of the sum order.

PROPOSITION 8.2 *Any inductive $*$ -semiring satisfies (11) and (13), i.e., the identities $1^*1^* = 1^*$ and $1^*(1^*a)^* = 1^*a^*$. Moreover, any inductive $*$ -semiring satisfies $1^*a \leq a1^*$.*

Proof. The identity $1^* = 1^*1^*$ holds by Theorem 8.1 and the description of the initial iteration semiring.

Now for the inequality $1^*a \leq a1^*$. We have $1(a1^*) + a = a(1^* + 1) = a1^*$. Thus, $1^*a \leq a1^*$.

Last for (13). On one hand, $a \leq 1^*a$, and thus $a^* \leq (1^*a)^*$ and $1^*a^* \leq 1^*(1^*a)^*$. On the other hand, $1(1^*(1^*a)^*) + a^* \leq 1^*(1^*a)^* + (1^*a)^* = (1^* + 1)(1^*a)^* = 1^*(1^*a)^*$, and thus $1^*a^* \leq 1^*(1^*a)^*$. $\square$

PROPOSITION 8.3 *In any symmetric inductive $*$ -semiring S , $1^*a = a1^*$ for all $a \in S$.*

Proof. We have seen that $1^*a \leq a1^*$. Since $(1^*a)1 + a = (1^* + 1)a = 1^*a$, it holds that $a1^* \leq 1^*a$. $\square$

Consider now $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ equipped with the *sum order*, denoted $\leq$. We claim that $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is a symmetric inductive $*$ -semiring. In [19], it is shown that if an ordered semiring equipped with a star operation is ordered by the sum order, then it is a symmetric inductive $*$ -semiring iff it satisfies $aa^* + 1 = a^*$ and

$$\begin{aligned} ax + b = x &\Rightarrow a^*b \leq x \\ xa + b = x &\Rightarrow ba^* \leq x. \end{aligned}$$

In order to prove these properties hold, we describe all solutions of a linear fixed point equation over $\mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle$.

PROPOSITION 8.4 *Let s, r be series in $\mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle$.*

1. *If s is proper, then the equation $x = sx + r$ has s^*r as its unique solution.*
2. *If $s = 1 + s_0$, where s_0 is proper, then the solutions of $x = sx + r$ are the series of the form $s^*r + 1^*s_0^+t + t$, where t is any series.*
3. *If $s = k + s_0$, where s_0 is proper and $k \in \mathbb{N}_\infty$, $k \neq 0, 1$, then the solutions of $x = sx + r$ are the series of the form $s^*r + 1^*s_0^*t = s^*(r + t)$, where t is any series.*

Proof. For the first claim, see [5]. Assume that $s = k + s_0$ where s_0 is proper and $k \in \mathbb{N}_\infty$, $k \neq 0$. Consider the equation $x = f(x)$, where $f(x) = sx + r$. Since $\mathbb{N}_\infty$ is continuous and $x \leq f(x)$, all solutions can be obtained by starting with a series t and forming the increasing sequence $f^n(t)$, for $n \geq 0$, and taking the supremum of this sequence. Since $f^n(t) = s^n t + s^{n-1}r + \dots + r$, this gives $s^*r + \sup_{n \geq 0} s^n t$. But for each n , $s^n t = (k + s_0)^n t$, and using the expansion $(k + s_0)^n = k^n + \binom{n}{1} k^{n-1} s_0 + \binom{n}{2} k^{n-2} s_0^2 + \dots + s_0^n$, we obtain that

$$\sup_{n \geq 0} s^n t = \begin{cases} 1^*s_0^+t + t & \text{if } k = 1 \\ 1^*s_0^*t & \text{if } k > 1. \end{cases}$$

$\square$

COROLLARY 8.5 *Let s, r be series in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ and consider the equation $x = sx + r$ with least solution s^*r . If z is any rational solution, then there is a rational series $p \in \mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ with $z = s^*r + p$.*

Proof. When s is proper, s^*r is the only solution and our claim is clear. Assume that $s = 1 + s_0$, where s_0 is proper (and rational). Then the least solution is $s^*r = 1^*s_0^*r$, and any other solution z is of the form $z = s^*r + 1^*s_0^+t + t = 1^*s_0^*r + 1^*s_0^+t + t$. We see that $z = s^*r + z$, so if z is rational, then it is the sum of the least solution with a rational series. The last case is when $s = k + s_0$, where s_0 is proper and $k \in \mathbb{N}_\infty$, $k > 1$. Then the least solution is $s^*r = 1^*s_0^*r$, and any other solution z is of the form $z = 1^*s_0^*r + 1^*s_0^*t$. We again have $z = s^*r + z$, so that if z is rational, then it is the sum of the least solution with a rational series. $\square$

Symmetrically, we have:

COROLLARY 8.6 *Let s, r be series in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ and consider the equation $x = xs + r$ with least solution rs^* . If z is any rational solution, then there is a rational series $p \in \mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ with $z = rs^* + p$.*

The main result of this section is:

THEOREM 8.7 *For each Σ , $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ (equipped with the sum order) is freely generated by Σ in the class of all inductive $*$ -semirings satisfying $a1^* \leq 1^*a$. In detail, for any inductive $*$ -semiring S satisfying $a1^* \leq 1^*a$ and for any function $h : \Sigma \rightarrow S$ there is a unique inductive $*$ -semiring morphism $h^\sharp : \mathbb{N}_\infty\langle\langle\Sigma^*\rangle\rangle \rightarrow S$ extending h .*

Proof. We have already proved that $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ equipped with the sum order is an inductive $*$ -semiring satisfying $a1^* \leq 1^*a$. Suppose that S and h are given. Since S satisfies $a1^* \leq 1^*a$, S is an iteration semiring in the variety $\mathcal{V}$ described in the previous section. By Theorem 7.14, h extends to a morphism $h^\sharp$ of $*$ -semirings. To see that $h^\sharp$ preserves the order, assume that $s, s' \in \mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ with $s \leq s'$. Then there exists a rational series r in $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ with $s + r = s'$. Since $h^\sharp$ preserves $+$, also $sh + rh = s'h$. But the order on S contains the sum order, so that $sh \leq s'h$. The fact that $h^\sharp$ is unique follows from Theorem 7.14. $\square$

COROLLARY 8.8 *For each Σ , $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the class of all symmetric inductive $*$ -semirings.*

COROLLARY 8.9 *For each Σ , $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is freely generated by Σ in the class of all sum ordered (symmetric) inductive $*$ -semirings.*

Let $k \geq 1$ and consider the $*$ -semiring $\mathbf{k}$. Equipped with the natural order, $\mathbf{k}$ is a continuous semiring and thus $\mathbf{k}\langle\langle\Sigma^*\rangle\rangle$ is also a continuous semiring and a symmetric inductive $*$ -semiring. We can write each series $s \in \mathbf{k}\langle\langle\Sigma^*\rangle\rangle$ in a unique way as the sum of series $s_0, s_1, \dots, s_k$, where each nonzero coefficient in any s_i is i . (Of course, s_0 is the 0 series.) Now by finiteness, it is known that s is rational iff each s_i is rational iff the support of each s_i is regular. Using this, we do not have the problem encountered in connection with the ordering of $\mathbb{N}_\infty^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$, the sum order and the pointwise order are equivalent on $\mathbf{k}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$. Using Theorem 7.19 we have:

THEOREM 8.10 *For each $k \geq 1$, $\mathbf{k}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle$ is both the free inductive $*$ -semiring and the free symmetric inductive $*$ -semiring on the set Σ satisfying the identity $k = k + 1$.*

Proof. Let S be an inductive $*$ -semiring satisfying $k = k + 1$. Since $1 \leq 1^*$ in S , also $k \leq 1^*$. Since $k + 1 = k$, also $1^* \leq k$. Thus $1^* = k$. By Theorem 7.19, every function $\Sigma \rightarrow S$ extends to a morphism $h^\sharp : \mathbf{k}^{\text{rat}}\langle\langle\Sigma^*\rangle\rangle \rightarrow S$ of iteration semirings. The extension $h^\sharp$ is monotone (and unique). $\square$

References

- [1] L. Aceto, Z. Ésik and A. Ingólfssdóttir, Equational theories of tropical semirings. Foundations of software science and computation structures (Genova, 2001). *Theoret. Comput. Sci.*, 298(2003), 417–469.
- [2] K.B. Arhangelsky and P.V. Gorshkov, Implicational axioms for the algebra of regular languages (in Russian). *Doklady Akad. Nauk, USSR, ser A.*, 10(1987), 67–69.

- [3] M.-P. Béal, S. Lombardy and J. Sakarovitch, On the equivalence of $\mathbb{Z}$ -automata, in: *ICALP 2005*, LNCS 3580, Springer, 2005, 397–409.
- [4] M.-P. Béal, S. Lombardy and J. Sakarovitch, Conjugacy and equivalence of weighted automata and functional transducers, in: *CSR 2006*, LNCS 3967, Springer, 2006, 58–69.
- [5] J. Berstel and Ch. Reutenauer, *Rational Series and Their Languages*, Springer, 1988.
- [6] S.L. Bloom and Z. Ésik, Matrix and matricial iteration theories, Part I, *J. Comput. Sys. Sci.*, 46(1993), 381–408.
- [7] S.L. Bloom and Z. Ésik, Equational axioms for regular sets, *Mathematical Structures in Computer Science*, 3(1993), 1–24.
- [8] S.L. Bloom and Z. Ésik, *Iteration Theories: The Equational Logic of Iterative Processes*, EATCS Monographs on Theoretical Computer Science, Springer-Verlag, 1993.
- [9] S.L. Bloom and Z. Ésik, Two axiomatizations of a star semiring quasi-variety, *EATCS Bulletin*, 59, June 1996, 150–152.
- [10] S.L. Bloom and Z. Ésik, The equational logic of fixed points, *Theoretical Computer Science*, 179(1997), 1–60.
- [11] S. L. Bloom, Z. Ésik and W. Kuich, Partial Conway and iteration semirings, *Fundamenta Informaticae*, to appear.
- [12] M. Boffa, A remark on complete systems of rational identities (French), *RAIRO Inform. Theor. Appl.*, 24(1990), 419–423.
- [13] M. Boffa, A condition implying all rational identities (French), *RAIRO Inform. Theor. Appl.*, 29(1995), 515–518.
- [14] A. Bonnier-Rigny and D. Krob, A complete system of identities for one-letter rational expressions with multiplicities in the tropical semiring, *Theoretical Computer Science*, 134(1994), 27–50.
- [15] J.C. Conway. *Regular Algebra and Finite Machines*, Chapman and Hall, London, 1971.
- [16] S. Eilenberg, *Automata, Languages, and Machines*. vol. A, Academic Press, 1974.
- [17] Z. Ésik, Identities in iterative and rational algebraic theories, *Computational Linguistics and Computer Languages*, XIV(1980), 183–207.
- [18] Z. Ésik, Group axioms for iteration, *Information and Computation*, 148(1999), 131–180.
- [19] Z. Ésik and W. Kuich, Inductive $*$ -semirings, *Theoret. Comput. Sci.*, 324(2004), 3–33.
- [20] J.S. Golan, *The Theory of Semirings with Applications in Computer Science*, Longman Scientific and Technical, 1993.
- [21] G. Grätzer, *Universal Algebra*, Springer, 1979.
- [22] D. Kozen, A completeness theorem for Kleene algebras and the algebra of regular events, *Technical Report*, Cornell University, Department of Computer Science, 1990.
- [23] D. Kozen, A completeness theorem for Kleene algebras and the algebra of regular events, *Inform. and Comput.*, 110(1994), 366–390.
- [24] D. Krob, Complete semirings and monoids (French), *Semigroup Forum*, 3(1987), 323–329.

- [25] D. Krob, Complete systems of B-rational identities, *Theoretical Computer Science*, 89(1991), 207–343.
- [26] D. Krob, Models of a K -rational identity system, *J. Computer and System Sciences*, 45(1992), 396–434.
- [27] D. Krob, The equality problem for rational series with multiplicities in the tropical semiring is undecidable, *International Journal of Algebra and Computation*, 4(1994), 405–425.
- [28] W. Kuich, The Kleene and Parikh theorem in complete semirings, in: *ICALP 1987*, LNCS 267, Springer, 1987, 212–225.
- [29] M. Morisaki and K. Sakai, A complete axiom system for rational sets with multiplicity, *Theoretical Computer Science*, 11(1980), 79–92.
- [30] V.N. Redko, On the determining totality of relations of an algebra of regular events (in Russian), *Ukrainian Math. Ž.*, 16(1964), 120–126.
- [31] V.N. Redko, On algebra of commutative events (in Russian), *Ukrainian Math. Ž.*, 16(1964), 185–195.
- [32] A. Salomaa, Two complete axiom systems for the algebra of regular events, *Journal of the Association for Computing Machinery*, 13(1966), 158–169.